

# UNIT 2

## *Chapter 2- Factor Theorem and Inequalities*

*WORKBOOK*

*MHF4U*

© MARK ANDERSON

WWW.ANDERTOONS.COM



"I don't like long division; I always feel bad  
for the remainders."

**W1 – 2.1 – Long Division of Polynomials and The Remainder Theorem**

**MHF4U**

**Jensen**

**1)** Use the remainder theorem to determine the remainder when  $2x^3 + 7x^2 - 8x + 3$  is divided by each binomial.

a)  $x + 1$

b)  $x - 2$

c)  $x + 3$

**2)a)** Divide  $x^3 + 3x^2 - 2x + 5$  by  $x + 1$ . Express the result in quotient form.

**b)** Write the corresponding statement that can be used to check the division.

**3)** Divide  $3x^4 - 4x^3 - 6x^2 + 17x - 8$  by  $3x - 4$ . Express the result in quotient form.

**b)** Write the corresponding statement that can be used to check the division.

**4)** Perform each division. Express the result in quotient form.

**a)**  $x^3 + 7x^2 - 3x + 4$  divided by  $x + 2$

**b)**  $6x^3 + x^2 - 14x - 6$  divided by  $3x + 2$

**c)**  $10x^3 + 11 - 9x^2 - 8x$  divided by  $5x - 2$

**d)**  $11x - 4x^4 - 7$  divided by  $x - 3$

e)  $6x^3 + x^2 + 7x + 3$  divided by  $3x + 2$

f)  $8x^3 + 4x^2 - 31$  divided by  $2x - 3$

g)  $6x^2 - 6 + 8x^3$  divided by  $4x - 3$

**5)** The volume, in cubic cm, of a rectangular box can be modelled by the polynomial expression  $2x^3 + 17x^2 + 38x + 15$ . Determine possible dimensions of the box if the height, in cm, is given by  $x + 5$ .

**6)** Determine the value of  $k$  such that when  $P(x) = kx^3 + 5x^2 - 2x + 3$  is divided by  $x + 1$ , the remainder is 7.

### ANSWER KEY

**1)** a) 16 b) 31 c) 36

**2)a)**  $\frac{x^3+3x^2-2x+5}{x+1} = x^2 + 2x - 4 + \frac{9}{x+1}$    **b)**  $x^3 + 3x^2 - 2x + 5 = (x + 1)(x^2 + 2x - 4) + 9$

**3)a)**  $\frac{3x^4-4x^3-6x^2+17x-8}{3x-4} = x^3 - 2x + 3 + \frac{4}{3x-4}$    **b)**  $3x^4 - 4x^3 - 6x^2 + 17x - 8 = (3x - 4)(x^3 - 2x + 3) + 4$

**4)a)**  $\frac{x^3+7x^2-3x+4}{x+2} = x^2 + 5x - 13 + \frac{30}{x+2}$    **b)**  $\frac{6x^3+x^2-14x-6}{3x+2} = 2x^2 - x - 4 + \frac{2}{3x+2}$

**c)**  $\frac{10x^3-9x^2-8x+11}{5x-2} = 2x^2 - x - 2 + \frac{7}{5x-2}$    **d)**  $\frac{-4x^4+11x-7}{x-3} = -4x^3 - 12x^2 - 36x - 97 - \frac{298}{x-3}$

**e)**  $\frac{6x^3+x^2+7x+3}{3x+2} = 2x^2 - x + 3 - \frac{3}{3x+2}$    **f)**  $\frac{8x^3+4x^2-31}{2x-3} = 4x^2 + 8x + 12 + \frac{5}{2x-3}$

**g)**  $\frac{6x^2-6+8x^3}{4x-3} = 2x^2 + 3x + \frac{9}{4} + \frac{3}{4(4x-3)}$

**5)**  $2x^3 + 17x^2 + 38x + 15 = (x + 5)(x + 3)(2x + 1)$

**6)**  $k = 3$

## W2 – 2.1 – Synthetic Division

MHF4U

Jensen

1) Calculate each of the following using synthetic division. Express your answer using the statement that could be used to check the division.

a)  $x^3 - 7x - 6$  divided by  $x - 3$

b)  $2x^3 - 7x^2 - 7x + 19$  divided by  $x - 1$

c)  $6x^4 + 13x^3 - 34x^2 - 47x + 28$  divided by  $x + 3$

d)  $2x^3 + x^2 - 22x + 20$  divided by  $2x - 3$

e)  $12x^4 - 56x^3 + 59x^2 + 9x - 18$  divided by  $2x + 1$

f)  $6x^3 - 15x^2 - 2x + 5$  divided by  $2x - 5$

**g)**  $x^3 - 2x + 1$  divided by  $x - 4$

**h)**  $x^3 + 2x^2 - 6x + 1$  divided by  $x + 2$

**2)** Divide  $x^4 - 16x^3 + 4x^2 + 10x - 11$  by each of the following binomials...

**a)**  $x - 2$

**b)**  $x + 4$

**3)** Are either of the binomials in question #2 factors of  $x^4 - 16x^3 + 4x^2 + 10x - 11$ ? Explain.

## ANSWER KEY

- 1)a)**  $x^3 - 7x - 6 = (x - 3)(x^2 + 3x + 2)$    **b)**  $2x^3 - 7x^2 - 7x + 19 = (x - 1)(2x^2 - 5x - 12) + 7$   
**c)**  $6x^4 + 13x^3 - 34x^2 - 47x + 28 = (x + 3)(6x^3 - 5x^2 - 19x + 10) - 2$   
**d)**  $2x^3 + x^2 - 22x + 20 = (2x - 3)(x^2 + 2x - 8) - 4$   
**e)**  $12x^4 - 56x^3 + 59x^2 + 9x - 18 = (2x + 1)(6x^3 - 31x^2 + 45x - 18)$    **f)**  $6x^3 - 15x^2 - 2x + 5 = (2x - 5)(3x^2 - 1)$   
**g)**  $x^3 - 2x + 1 = (x - 4)(x^2 + 4x + 14) + 57$    **h)**  $x^3 + 2x^2 - 6x + 1 = (x + 2)(x^2 - 6) + 13$   
**2)a)**  $x^4 - 16x^3 + 4x^2 + 10x - 11 = (x - 2)(x^3 - 14x^2 - 24x - 38) - 87$   
**b)**  $x^4 - 16x^3 + 4x^2 + 10x - 11 = (x + 4)(x^3 - 20x^2 + 84x - 326) + 1293$   
**3)** No, because for each division problem, there is a remainder.

**W3 – 2.2 – Factor Theorem**

**MHF4U**

*Jensen*

**1)** Determine if  $x + 3$  is a factor of each polynomial:

a)  $x^3 + x^2 - x + 6$

b)  $2x^3 + 9x^2 + 10x + 3$

c)  $x^3 + 27$

**2)** Find possible factors of the following polynomials using integral zero theorem. Then, factor the polynomial.

a)  $x^3 + 3x^2 - 6x - 8$

b)  $x^3 + 4x^2 - 15x - 18$

c)  $x^3 - 3x^2 - 10x + 24$

**3)** Factor by grouping:

a)  $x^3 + x^2 - 9x - 9$

b)  $2x^3 - x^2 - 72x + 36$

**4)** Determine a value of k so that  $x+2$  is a factor of  $x^3 - 2kx^2 + 6x - 4$ .

**5)** Find possible factors of the following polynomials using integral zero theorem. Then, factor the polynomial.

**a)**  $3x^3 + x^2 - 22x - 24$

**b)**  $2x^3 - 9x^2 + 10x - 3$

**c)**  $6x^3 - 11x^2 - 26x + 15$

**d)**  $4x^3 + 3x^2 - 4x - 3$

**6) Factor each polynomial**

a)  $2x^3 + 5x^2 - x - 6$

b)  $4x^3 - 7x - 3$

c)  $x^4 - 15x^2 - 10x + 24$

## ANSWER KEY

**1)a) No b) Yes c) Yes**

**2)a)  $(x - 2)(x + 1)(x + 4)$  b)  $(x - 3)(x + 1)(x + 6)$  c)  $(x - 4)(x - 2)(x + 3)$**

**3)a)  $(x - 3)(x + 1)(x + 3)$  b)  $(x - 6)(x + 6)(2x - 1)$**

**4)  $k = -3$**

**5)a)  $(x - 3)(x + 2)(3x + 4)$  b)  $(x - 3)(x - 1)(2x - 1)$  c)  $(x - 3)(2x - 1)(3x + 5)$  d)  $(x - 1)(x + 1)(4x + 3)$**

**6)a)  $(x - 1)(x + 2)(2x + 3)$  b)  $(x + 1)(2x - 3)(2x + 1)$  c)  $(x - 4)(x - 1)(x + 2)(x + 3)$**

## W4 – 2.3 – Solving Polynomial Equations

MHF4U

Jensen

1) Determine the solutions of the following polynomials.

a)  $(3x + 2)(x + 9)(x - 2) = 0$

b)  $(x^2 + 1)(x - 4) = 0$

2) Determine the solutions of the following polynomials by factoring. Use the tools you have learned this unit to help you. (remainder theorem, integral zero theorem, division etc.)

a)  $x^3 - 4x^2 - 3x + 18 = 0$

b)  $x^3 - 3x^2 - 4x + 12 = 0$

c)  $x^4 - x^3 - 11x^2 + 9x + 18 = 0$

d)  $x^3 - 64 = 0$

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

e)  $2x^3 - 7x^2 + 10x - 5 = 0$

**3)** Solve each equation by first factoring the sum or difference of cubes.

**a)**  $x^3 - 8 = 0$

**b)**  $x^3 + 27 = 0$

**4)** Solve by factoring

**a)**  $x^3 - 4x^2 - 7x + 10 = 0$

**b)**  $2x^3 - 11x^2 + 12x + 9 = 0$

**c)**  $x^4 - x^3 - 2x - 4 = 0$

### ANSWER KEY

**1a)**  $\left(-\frac{2}{3}, 0\right), (-9, 0), (2, 0)$     **b)**  $(4, 0)$

**2a)**  $(-2, 0)$  and  $(3, 0)$     **b)**  $(3, 0), (-2, 0), (2, 0)$     **c)**  $(-1, 0), (2, 0), (-3, 0), (3, 0)$     **d)**  $(4, 0)$     **e)**  $(1, 0)$

**3)a)**  $(2, 0)$     **b)**  $(-3, 0)$

**4)a)**  $(5, 0), (-2, 0), (1, 0)$     **b)**  $(-0.5, 0)$  and  $(3, 0)$     **c)**  $(-1, 0)$  and  $(2, 0)$

## W5 – 2.4 – Families of Polynomial Functions

MHF4U

Jensen

1) The zeros of a quadratic function are -7 and -3.

a) Determine an equation for the family of quadratic functions with these zeros.





b) Write equations for two functions that belong to this family.

c) Determine an equation for the member of the family that passes through the point (2, 18).

2) Examine the following functions. Which function does not belong to the same family?

- a)  $y = 1.5(x + 4)(x - 5)(x - 2)$
- b)  $y = -1.5(x - 2)(x - 5)(x + 4)$
- c)  $y = 1.5(x - 2)(x + 4)(x - 2)$
- d)  $y = 3(x - 5)(x - 2)(x + 4)$

3) The graphs of four polynomial functions are given. Which graphs represent functions that belong to the same family?




**4)a)** Determine an equation for the family of cubic functions with zeros  $-2$ ,  $-1$ , and  $\frac{1}{2}$

**b)** Write equations for two functions that belong to this family.

**c)** Determine an equation for the member of the family whose graph has a  $y$ -intercept of  $6$ .

**d)** Sketch a graph of the function from part c).





**5)a)** Determine an equation for the family of cubic functions with zeros  $1 \pm \sqrt{2}$  and  $-\frac{1}{2}$

**b)** Determine an equation for the member of the family whose graph passes through the point  $(3, 35)$ .

**6)a)** Determine an equation for the family of quartic functions with zeros 3 (order 2) and  $-4 \pm \sqrt{3}$ .

**b)** Determine an equation for the member of the family whose graph passes through the point  $(1, -22)$ .

7) Determine an equation for each of the following functions



### ANSWER KEY

- 1)a)  $y = k(x + 7)(x + 3)$    b) answer will vary   c)  $y = \frac{2}{5}(x + 7)(x + 3)$       2) C      3) A, B, D  
 4)a)  $y = k(x + 2)(x + 1)(2x - 1)$    b) answer will vary   c)  $y = -3(x + 2)(x + 1)(2x - 1)$    d) see posted  
 5)a)  $y = k(x^2 - 2x - 1)(2x + 1)$    b)  $y = \frac{5}{2}(x^2 - 2x - 1)(2x + 1)$   
 6)a)  $y = k(x - 3)^2(x^2 + 8x + 13)$    b)  $y = -\frac{1}{4}(x - 3)^2(x^2 + 8x + 13)$   
 7)a)  $y = -2(x + 2)(x - 1)(x - 3)$    b)  $y = (x + 3)^2(x - 1)(2x - 3)$

## W6 – 2.5 – Solving Inequalities

MHF4U

Jensen

1) Solve each linear inequality

a)  $x + 3 \leq 5$

b)  $7x < 4 + 3x$

2) Solve each inequality by graphing

a)  $(x + 3)(x - 2) > 0$

b)  $(x + 2)(3 - x)(x + 1) < 0$

**3)** Solve each of the following polynomial inequalities

**a)**  $x^2 - 7x + 10 \geq 0$

**b)**  $x^3 + 6x^2 - 16x > 0$

**c)**  $-x^2 + 36 \geq 0$

**d)**  $x^4 - 26x^2 + 25 > 0$

$$\mathbf{e)} \quad x^3 - 3x^2 \geq 25x - 75$$

$$\mathbf{f)} \quad -x^3 + 28x + 48 \geq 0$$

$$\mathbf{g)} \quad x^3 - 2x^2 - 5x + 6 < 0$$

$$\mathbf{h)} \quad 5x^3 - 12x^2 - 11x + 6 \leq 0$$

- 4)** The price,  $p$ , in dollars, of a stock  $t$  years after 1999 can be modelled by the function  $p(t) = 0.5t^3 - 5.5t^2 + 14t$ . When will the stock be more than \$90? You may use technology to help you determine the solution.

### ANSWER KEY

- 1)a)**  $x \leq 2$    **b)**  $x < 1$   
**2)a)**  $x < -3$  or  $x > 2$    **b)**  $-2 < x < -1$  or  $x > 3$   
**3)a)**  $x \leq 2$  or  $x \geq 5$    **b)**  $-8 < x < 0$  or  $x > 2$    **c)**  $-6 \leq x \leq 6$    **d)**  $x < -5$  or  $-1 < x < 1$  or  $x > 5$   
**e)**  $-5 \leq x \leq 3$  or  $x \geq 5$    **f)**  $x \leq -4$  or  $-2 \leq x \leq 6$    **g)**  $x < -2$  or  $1 < x < 3$   
**h)**  $x \leq -1$  or  $\frac{2}{5} < x < 3$   
**4)** after 10 years (2009)