
# Unit 3 Pre-Test Review – Exponential and Logarithmic Functions

MHF4U

Jensen


1) Sketch a graph of each function. Then, sketch a graph of the inverse of each function. Label each graph with its equation. Also, complete the table of information for each function

**a)** 
$$f(x) = 2^x$$



f(x) =	$f^{-1}(x) =$
<i>x</i> -int:	x-int:
y-int:	<i>y</i> -int:
Domain:	Domain:
Range:	Range:
Asymptote:	Asymptote:

**b)** 
$$g(x) = \left(\frac{1}{4}\right)^x$$




g(x) =	$g^{-1}(x) =$
x-int:	<i>x</i> -int:
y-int:	<i>y</i> -int:
Domain:	Domain:
Range:	Range:
Asymptote:	Asymptote:


2) State the domain and range for the function, shown below.

Domain:


Range:




A)




B)




C)



**Graph:** 





x 0 2

**Equation:** 

Letter of Graph of Inverse:

**Equation of inverse:** 

**4)** Rewrite each equation in logarithmic form.

a) 
$$4^3 = 64$$

**b)** 
$$28 = 3^x$$

**c)** 
$$6^3 = y$$

**d)** 
$$512 = 2^9$$

5) Rewrite each equation in exponential form.

**a)** 
$$7 = \log_2 128$$

**b)** 
$$x = \log_b n$$

**c)** 
$$5 = \log_3 243$$
 **d)**  $19 = \log_b 4$ 

**d)** 
$$19 = \log_b 4$$

- **6)** Evaluate without a calculator. Show your work.
- a)  $\log_2 16$

**b)**  $\log_3 81$ 

Use either:

**Rule:** if  $x^a = x^b$ , then a = b

**Rule:**  $\log_a(a^b) = b$ 

c)  $\log_4\left(\frac{1}{16}\right)$ 

**d)** log 0.000 001

- 7) Evaluate each of the following without a calculator using the power law of logarithms.
- **a)**  $\log_2 32^3$
- **b)**  $\log 1000^{-2}$
- c)  $\log 0.001^{-1}$
- **d)**  $\log_{\frac{1}{4}} \left(\frac{1}{16}\right)^4$

**8)** Solve for x, correct to 3 decimal places.

**a)** 
$$x = \log_3 17$$

**b)** 
$$\log_2 0.35 = x$$
 **c)**  $4^x = 10$ 

c) 
$$4^x = 10$$

**d)** 
$$80 = 100 \left(\frac{1}{2}\right)^x$$

0)	-£   £		Da	حججات احجدته جام
9) Use the change	or base for	mula to evaluate	. Rouna to one	e decimai piace.

**a)** log<sub>9</sub> 12

**b)**  $\log_{0.25} 52$ 

10) Write as a single logarithm. Then evaluate without a calculator.

a)  $\frac{\log 16}{\log 4}$ 

**b)**  $\frac{\log\left(\frac{8}{27}\right)}{\log\left(\frac{2}{3}\right)}$ 

11) Solve, to two decimal places

**a)**  $\log 4^x = 7$ 

**b)**  $12 = \log_3 4^m$ 

- **12)** An investment earns 12% interest, compounded annually. The amount, A, that he investment is worth as a function of time, t, in years, is given by  $A = 1500(1.12)^t$ . Use the equation to determine...
- a) the value of the investment after 4 years

**b)** how long it will take for the investment to double in value

13) Write as a single logarithm

a) 
$$\log_7 8 + \log_7 4 - \log_7 16$$

**b)** 
$$2 \log a + \log(3b) - \frac{1}{2} \log c$$

14) Write as a sum or difference of logarithms. Simplify if possible.

a) 
$$\log(a^2bc)$$

**b)** 
$$\log\left(\frac{k}{\sqrt{m}}\right)$$

- **15)** Evaluate, using the laws of logarithms.
- **a)**  $\log_6 8 + \log_6 27$

**b)**  $\log_4 128 - \log_4 8$ 

c)  $2 \log 2 + 2 \log 5$ 

**d)**  $2 \log 3 + \log \left(\frac{25}{2}\right)$ 

- 16) Simplify
- a)  $\log(2m+6) \log(m^2-9)$

**b)**  $\log(x^2 + 2x - 15) - \log(x^2 - 7x + 12)$ 

- 17) Write each as a power of 4
- **a)** 64

**b)**  $\frac{1}{16}$ 

c)  $(\sqrt[3]{8})^5$ 

**18)** Write 20 as a power of 5.

19) Solve each equation

a) 
$$3^{5x} = 27^{x-1}$$

**b)** 
$$8^{2x+1} = 32^{x-1}$$

**20)** Solve exactly. Then use your calculator to evaluate correct to 3 decimal places.

**a)** 
$$3^{x-2} = 5^x$$

**b)** 
$$2^{k-2} = 3^{k+1}$$

21) Solve the following equations; round to 2 decimal places where appropriate.

**a)** 
$$3^x = 12$$

**b)** 
$$10 = 2 \cdot 4^{x+2}$$

c) 
$$3^x = 4^{1-x}$$

**22)** Solve each equation. Check for extraneous routes.

a) 
$$4^{2x} - 4^x - 20 = 0$$

**b)** 
$$2^x + 12(2)^{-x} = 7$$

## 23) Solve each equation

**a)** 
$$\log_4 x = 1.8$$

**b)** 
$$log_5 x - log_5 (x - 2) = 1$$

**c)** 
$$5^{2x} = 2(5)^x + 1$$

**24)** Solve

**a)** 
$$\log(2x + 10) = 2$$

**b)** 
$$1 - \log(2x) = 0$$

25) Solve. Check for extraneous roots.

**a)** 
$$\log_2 x + \log_2 (x+2) = 3$$

**b)** 
$$\log_3(3x + 7) = 2$$

c) 
$$\log_5(2x+1) = 1 - \log_5(x+2)$$

#### Section 6: 7.4 - Applications

### **Exponential Formulas**

$$A(t) = A_0 (1+i)^t$$

 $A(t) = A_0 \left(\frac{1}{2}\right)^{\frac{t}{H}}$ 

$$A(t) = A_0(2)^{\frac{t}{D}}$$

general, where *i* is percent growth(+) or decay(-)

half-life, H is the half-life period

doubling, *D* is the doubling period

#### **Logarithmic Formulas**

$$pH = -\log[H^+]$$

 $\beta_2 - \beta_1 = 10 \log \left(\frac{I_2}{I_1}\right)$ 

 $M = \log(\frac{I}{I_0})$ 

Where pH is acidity and [H+] is concentration of hydronium ions mol/L

Where  $\beta$  is loudness in dB and I is intensity of sound in W/m<sup>2</sup>

Where M is magnitude measure by richters, I is intensity

- **26)** When you drink a cup of coffee or a glass of cola, or when you eat a chocolate bar, the percent, P, of caffeine remaining in your bloodstream is related to the elapsed time, t, in hours by  $t = 5\left(\frac{\log P}{\log 0.5}\right)$
- a) How long will it take for the amount of caffeine to drop to 20% of the amount consumed?

**b)** Suppose you drink a cup of coffee at 9:00 am, what percent of the caffeine will remain in your body at noon?


27) A 50-mg sample of cobalt-60 decays to 40 mg after 1.6	minutes.
a) Determine the half-life of cobalt-60.	
<b>b)</b> How long will it take for the sample to decay to 5% of its	initial amount?
28) Determine the pH, correct to one decimal place, of a sc	olution with each hydronium ion concentration.
a) 0.000 316 mol/L	<b>b)</b> 7.9 × 10 <sup>-9</sup> mol/L
29) Calculate the hydronium ion concentration, correct to twa) 2.2	vo decimal places, if the pH of a solution is <b>b)</b> 11.6
<b>30)</b> Use the sound level scale in your notes to answer the fo	ollowing:
a) How many times as intense is a normal conversation con	npared to a whisper?
<b>b)</b> How many times as intense is normal city traffic compar	ed to a shout?

<b>31)</b> The intensity of sound in a library is estimated to be one thousandth that of normal conversation. What is the decibel rating for the library?
<b>32)</b> How many times as intense is an earthquake with a magnitude of 7.2 than an earthquake with a magnitude of 5.6?
<b>33)</b> If an earthquake is 390 times as intense as an earthquake with a magnitude of 4.2 on the Richter scale, what is the magnitude of the more intense earthquake?
<b>34)</b> The absolute magnitude of star A is −4.5 and that of star B is 0.2. How many times as bright is star A than star B, to the nearest unit?


<b>35)</b> An altimeter is a device that measures the height of a plane above the ground. It works based on air pressure according to the formula $h=18400log\frac{P_0}{P}$ , where h is the height above the ground in metres, P is the air pressure at that height, and $P_0$ was the air pressure on the ground at takeoff. Air pressure is measure in kilopascals (kPa).
a) Air pressure on the ground was 102 kPa. If the airplane instruments measure a pressure of 32.5 kPa outside the plane, what is the height of the airplane to the nearest metre?
b) What is the outside air pressure for a plane flying at 11 000 metres? Assume a ground pressure 102.5 kPa. Round to one decimal place.
c) How high would a plane have to be flying when it encountered air pressure in the air that was half the air pressure on the ground? Round to the nearest meter.

**36)** Sketch a graph of each of the following exponential/logarithmic functions by applying transformations to the parent function. Make sure to identify key points such as asymptotes and x-intercepts.


**a)** 
$$f(x) = 2(2)^{-2x-2} + 1$$




**b)** 
$$g(x) = -(e)^{\frac{1}{2}(x+2)} + 4$$



c) 
$$h(x) = -2\log(x+4) + 1$$



**d)** 
$$j(x) = 3 \ln(x - 1) + 3$$



**37)** Solve each equation. Round your answer to 4 decimal places if necessary.

**a)** 
$$e^{3x} = 87$$

**b)** 
$$2e^{3x+1} = 70$$

$$\mathbf{c})\ln(x+1) = \ln(2x-5)$$

**d)** 
$$5 \ln x + 2 \ln x - 3 = 12$$

**e)** 
$$\ln(3x) = 2$$

$$f) 1 - 2e^{2x} = -19$$

#### **Answer Key**

See posted solutions for #1-3

**4)a)** 
$$\log_4 64 = 3$$
 **b)**  $\log_3 28 = x$  **c)**  $\log_6 y = 3$  **d)**  $\log_2 512 = 9$ 

**5)a)** 
$$2^7 = 128$$
 **b)**  $b^x = n$  **c)**  $3^5 = 243$  **d)**  $b^{19} = 4$ 

**10)a)** 
$$\log_4 16 = 2$$
 **b)**  $\log_{\frac{2}{3}} \left( \frac{8}{27} \right) = 3$ 

**13)a)** 
$$\log_7 2$$
 **b)**  $\log \left( \frac{3a^2b}{\sqrt{c}} \right)$ 

**14)a)** 
$$2 \log a + \log b + \log c$$
 **b)**  $\log k - \frac{1}{2} \log m$ 

**16)a)** 
$$\log \left( \frac{2}{m-3} \right)$$
 **b)**  $\log \left( \frac{x+5}{x-4} \right)$ 

17)a) 
$$4^3$$
 b)  $4^{-2}$  c)  $4^{\frac{5}{2}}$ 

18) 
$$5^{\frac{\log 20}{\log 5}}$$

**19)a)** 
$$x = -\frac{3}{2}$$
 b)  $x = -8$ 

**20)a)** 
$$x = \frac{2 \log 3}{\log 3 - \log 5} \cong -4.301$$
 **b)**  $k = \frac{2 \log 2 + \log 3}{\log 2 - \log 3} \cong -6.129$ 

**22)a)** 
$$x = \frac{\log 5}{\log 4} \cong 1.16$$
 **b)**  $x = 2$  or  $x = \frac{\log 3}{\log 2} \cong 1.58$ 

**23)a)** 12.13 **b)** 2.5 **c)** 
$$x = 0.548$$

**25)a)** 2 **b)** 
$$\frac{2}{3}$$
 **c)**  $\frac{1}{2}$ 

**29) a)** 
$$6.31 \times 10^{-3}$$
 mol/L **b)**  $2.51 \times 10^{-12}$  mol/L